

Missouri Pharmacy Program – Preferred Drug List



Onychomycosis Antifungal Agents Effective 12/07/05 Revised 07/03/2008

Preferred Agents

- Gris-Peg®
- Griseofulvin Susp
- Grifulvin® V Tablets
- Griseofulvin Tablets
- Griseofulvin Powder
- Terbinafine
- Ciclopirox Solution

Non-Preferred Agents

- Sporanox® Capsules
- Sporanox® Solution
- Itraconazole
- Fulvicin® Tablets
- Penlac®
- Grifulvin V Suspension
- Lamisil®
- CNL-8 Nail Kit

Approval Criteria	Denial Criteria	
Documented diagnosis of proximal or distal, white, subungual onychomycosis, identified through: o KOH microscopic exam o Fungal Culture, or o Nail Biopsy	Lack of adequate trial on required preferred agents	
> 30% nail plate involvement	Therapy will be denied if no approval criteria are met	
Failure to achieve desired therapeutic outcomes with trial on 1 preferred agents o Documented trial period for preferred agents o Documented ADE/ADR to preferred agents	Sporanox Only: Left ventricular dysfunction, such as congestive heart failure (CHF)	
Documented compliance on current therapy regimen		
See Appendix for maximum approvable durations of therapy	Drug Prior Authorization Hotline: (800) 392-8030	

Appendix

Product	Strength/Dose	Duration of Therapy	Anatomic location of infection
Lamisil	250 mg Once Daily	6 weeks	Fingernails
Lamisil	250 mg Once Daily	12 Weeks	Toenails
Sporanox	200 mg BID	1 week (3 weeks no therapy) for 3 cycles (Pulse)	Fingernails
Sporanox	200 mg Once Daily	12 weeks (or Pulse)	Toenails